
# IceBase: A suborbital survey to map geothermal heat flux under an ice sheet

Michael E. Purucker, Alan P.M. Vaughan, Jurgen Matzka, Nils Olsen, John Jorgensen, Chris Finlay, T. Rasmussen, Catherine Fox Maule, Jack Connerney, Brad Nelson, Richard Blakely, Robert Bracken, Sophie Nowicki, Guan Le, Terrence Sabaka, Todd Bonalsky, Weijia Kuang, D. Ravat, C. Ritz, Claire Bouligand, Carmen Gaina, Suzanne McEnroe, Vincent Lesur, Robert Tyler, Jared Espley, and Erwan Thebault






# A joint US-Danish effort

# Front

#### Magnetic crustal thickness/heat flux: Back (peelable)



### Plan for analysis of science data

- Remove airplane fields
- remove time-variable core field
- remove time-variable magnetospheric field
- remove tidal oceanic magnetic field
- remove ionospheric currents

remove coastal effect

- · level flight lines
- grid data
- Remove crustal field signatures unrelated to crustal thickness
- Determine depth to Curie isotherm using MCT (magnetic crustal thickness) and DBT (power-spectral) approaches.
- determine heat flux using 1-D heat conduction eq. or utilizing the four layer model (sediment, upper, middle, lower crust) of Fox Maule et al. (2009), with calibration/validation provided by other independent
- Ice sheet evolution using these boundary conditions.

# Science concept

Fox Maule, Purucker, Olsen, Mosegaard, 2005, Science, Heat Flux Anomalies in Antarctica revealed by satellite magnetic data

Summary: NASA will solicit suborbital missions as part of its Earth

suborbital magnetic survey of Greenland using NASA's Global Hawk

Venture program element in the coming year. These missions are

designed as complete PI-led investigations to conduct innovative

hypothesis or scientific question-driven approaches to pressing

questions in Earth System science. We propose to carry out a

unmanned aerial vehicle to produce the first-ever map of the

geothermal heat flux under an ice sheet. Better constraints on

geothermal heat flux will reduce the uncertainty in future sea level

rise, in turn allowing a more informed assessment of its impact on

mantle heat flux, and the tectonic history and heat production of the

geothermal heat flux influences the basal ice. Therefore heat flux is

magnetic data to constrain heat flux is possible because the magnetic

Curie temperature. The technique has applications to understanding

properties of rocks are temperature dependent until they reach the

the response of Greenland ice sheet to climate forcing because the

Greenland should be found in areas of very low heat flux, and the

identification of those areas is provided by this technique. Ice cores

from the areas of oldest ice help to decipher past temperatures and

Greenland Basal Heat Flux) is based on low-resolution satellite

observations collected by the CHAMP satellite between 2000 and

a factor of two, from 400 km resolution to approximately 250 km

2010. Those observations will be enhanced by the upcoming Swarm

gradient satellite mission, but the resolution will improve by less than

resolution. A high altitude, suborbital magnetic survey of Greenland

region associated with the Iceland mantle plume in E /SE Greenland,

Canadian Shield. Magnetic field measurements from 20 km altitude

are strongly preferred over lower altitude observations because of

calibration with sufficient sensitivity, and suppress local remanent

assessing the possible contributions from remanent magnetism and

variable susceptibility, and from other lithospheric processes such as

structure, volcanism and impact, from unmodeled external magnetic

magnetic field signatures. We validate our heat flux estimates by

fields, and from the assumptions utilized in the heat flux model.

their ability to sample the longest wavelengths, provide uniform

would provide a heat flux model with resolution comparable to the

crustal thickness, and would provide details of the high heat flux

and the low heat flux region in NW Greenland, adjacent to the

CO<sub>2</sub> contents. Our latest model of the geothermal heat flux under the

basal heat flux provides one of the boundary conditions. The


technique also helps to locate the oldest ice. The oldest ice in

Greenland ice sheet (http://websrv.cs.umt.edu/isis/index.php/

society. The geothermal heat flux depends on conditions such as

crust, all of which vary spatially. Underneath ice sheets, the

an important boundary condition in ice sheet modeling. Using




Moho assumed to coincide with Curie

isotherm and to be at 580 C


Magnetic field dF ~ Crustal thickness x magnetic susceptibility

Heat flux a ~ 1/Crustal thickness [Steady state 1-D heat conduction eq. w no lateral variations of material properties or heat production. 30-60% or more of surface heat flux is due to crustal heat production.]

# Basal boundary condition







v, SGT at Planetary Geodynamics Lab, GSFC/NASA, Greenbelt, MD 20771 For further information contact: Michael.E.Purucker@nasa.go